

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07D 401/12, 213/68, 213/89 A61K 31/44	A1	(11) International Publication Number: WO 90/06925 (43) International Publication Date: 28 June 1990 (28.06.90)
(21) International Application Number: PCT/SE89/00740 (22) International Filing Date: 20 December 1989 (20.12.89) (30) Priority data: 8804629-7 22 December 1988 (22.12.88) SE (71) Applicant: AKTIEBOLAGET HÄSSLE [SE/SE]; S-431 83 Mölndal (SE). (72) Inventors: BRÄNDSTRÖM, Arne, Elof ; Anders Mattssonsgatan 13 B, S-415 06 Göteborg (SE). LINDBERG, Per, Lennart ; Knapehall 64, S-436 39 Askim (SE). SUNDÉN, Gunnel, Elisabeth ; Frigångsgatan 10, S-413 01 Göteborg (SE).		(74) Agents: MIKSCHKE, Gerhard et al.; AB Astra, Patent Department, S-151 85 Södertälje (SE). (81) Designated States: AT, AT (European patent), AU, BE, BE (European patent), BF, (OAPI patent), BG, BJ (OAPI patent), BR, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CM (OAPI patent), DE, DE (European patent), DK, ES (European patent), FI, FR (European patent), GA (OAPI patent), GB, GB (European patent), HU, IT (European patent), JP, KP, KR, LK, LU, LU (European patent), MC, MG, ML (OAPI patent), MR (OAPI patent), MW, NL, NL (European patent), NO, RO, SD, SE, SE (European patent), SN (OAPI patent), SU, TD (OAPI patent), TG (OAPI patent). Published <i>With international search report.</i>
(54) Title: NEW THERAPEUTICALLY ACTIVE COMPOUND AND A PROCESS FOR ITS PREPARATION		
(57) Abstract The novel compound 5-fluoro-2[[[(4-cyclopropylmethoxy-2-pyridinyl)methyl]sulfinyl]-1H-benzimidazole and physiologically acceptable salts thereof as well as intermediates, pharmaceutical compositions containing the compound as active ingredient, and the use of the compound in medicine.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

New Therapeutically Active Compound and a Process for its Preparation

DESCRIPTION

5

Field of the invention

The object of the present invention is to provide a novel compound, and therapeutically acceptable salts thereof, which inhibit exogenously or endogenously stimulated gastric acid secretion and thus can be used in the prevention and treatment of peptic ulcer.

The present invention also relates to the use of the compound of the invention, especially therapeutically acceptable salts thereof, for inhibiting gastric acid secretion in mammals including man. In a more general sense, the compound of the invention may be used for prevention and treatment of gastrointestinal inflammatory diseases, and gastric acid-related diseases in mammals including man, such as gastritis, gastric ulcer, duodenal ulcer, reflux esophagitis, and Zollinger-Ellison syndrome. Furthermore, the compound may be used for treatment of other gastrointestinal disorders where gastric antisecretory effect is desirable e.g. in patients with gastrinomas, and in patients with acute upper gastrointestinal bleeding. It may also be used in patients in intensive care situations, and pre- and postoperatively to prevent acid aspiration and stress ulceration. The compound of the invention may also be used for treatment or prophylaxis of inflammatory conditions in mammals, including man, especially those involving lysozymal enzymes. Conditions that may be specifically mentioned are rheumatoid arthritis and gout. The compound may also be useful in the treatment of diseases related to bone metabolism disorders as well as the treatment of glaucoma.

The invention also relates to pharmaceutical compositions containing the compound of the invention, or a therapeutically acceptable salt thereof, as active ingredient. In a further aspect, the invention relates to processes for preparation of such new compound, to novel intermediates in the preparation of the compound of the invention, and to the use of the active compound for the preparation of pharmaceutical compositions for the medical use indicated above.

10

It is a specific primary object of the invention to provide a compound with a high level of bioavailability. The compound of the invention will also exhibit high stability properties at neutral pH and a high potency in regard to inhibition of gastric acid secretion. Bioavailability is defined as the fraction, or percent, of the administered dose of compound that is absorbed unchanged into the systemic blood. Potency is in this application defined as the ED₅₀ value.

20

Prior art and background of the invention

Benzimidazole derivatives intended for inhibiting gastric acid secretion are disclosed in numerous patent documents. Among these can be mentioned GB 1 500 043, GB 1 525 958, US 4 182 766, US 4 255 431, US 4 599 347, EP 124 495, US 4 555 518, US 4 727 150, US 4 628 098, EP 208 452 and Derwent abstract 87-294449/42. Benzimidazole derivatives proposed for use in the treatment or prevention of special gastrointestinal inflammatory diseases are disclosed in US 4 539 465.

30

The invention

Compounds described in the prior art, as described above, are effective acid secretion inhibitors, and are thus

35

useful as antiulcer compounds. In order to further enhance the usefulness of this type of drugs, a higher bioavailability has been desired, but still the compounds should have a high potency in inhibiting gastric acid secretion and also a high chemical stability at neutral pH.

It has been recognized that 2-[(2-pyridinylmethyl)-sulfinyl]-1H-benzimidazoles tested show a great variability in bioavailability as well as in potency and stability, and it is difficult to identify compounds possessing all the three advantageous properties. There is no guidance in the prior art on how to obtain compounds with this combination of properties.

It has been found that the compound of the invention shows exceedingly high bioavailability, and still the compound is very effective as inhibitor of gastric acid secretion and exhibits a high chemical stability in solution at a neutral pH. Thus the compound of the invention can be used in the indications given above in mammals including man.

The compound of the invention is 5-fluoro-2-[[[4-cyclopropylmethoxy-2-pyridinyl)methyl]sulfinyl])-1H-benzimidazole (compound I) and physiologically acceptable salts thereof. The compound of the invention has an asymmetric centre in the sulfur atom, i.e. exists as two optical isomers (enantiomers). Both the pure enantiomers, racemic mixtures (50% of each enantiomer) and unequal mixtures of the two are within the scope of the present invention. Also five synthetic intermediates and process for the preparation are within the scope.

Preparation

The compound of the invention, may be prepared according to the following method:

5 Oxidizing 5-fluoro-2[[[4-cyclopropylmethoxy-2-pyridinyl)methyl]-thio-1H-benzimidazole (compound II) to give the compound of the invention. This oxidation may be carried out by using an oxidizing agent such as nitric
10 acid, hydrogen peroxide, (optionally in the presence of vanadium compounds), peracids, peresters, ozone, dinitrogen tetroxide, iodosobenzene, N-halosuccinimide, 1-chlorobenzotriazole, t-butylhypochlorite, diazabicyclo-[2,2,2]-octane bromine complex, sodium metaperiodate,
15 selenium dioxide, manganese dioxide, chromic acid, ceric ammonium nitrate, bromine, chlorine, and sulfuric chloride. The oxidation usually takes place in a solvent such as halogenated hydrocarbons, alcohols, ethers, ketones.

20 The oxidation may also be carried out enzymatically by using an oxidizing enzyme or microbially by using a suitable microorganism.

25 Depending on the process conditions and the starting materials, the compound of the invention is obtained either in neutral or salt form. Both the neutral compound and the salts of this are included within the scope of the invention. Thus, basic, neutral or mixed salts may be
30 obtained as well as hemi, mono, sesqui or polyhydrates.

Alkaline salts of the compound of the invention are exemplified by its salts with Li^+ , Na^+ , K^+ , Mg^{2+} , Ca^{2+} , and $\text{N}^+(\text{R})_4$, where R is (1-4 C)alkyl. Particularly
35 preferred are the Na^+ , Ca^{2+} and Mg^{2+} salts. Especially preferred are the Na^+ and Mg^{2+} salts. Such salts may be

prepared by reacting the compound with a base capable of releasing the desired cation.

Examples of bases capable of releasing such cations, and
5 examples of reaction conditions are given below.

a) Salts wherein the cation is Li^+ , Na^+ or K^+ are prepared by treating the compound of the invention with LiOH , NaOH or KOH in an aqueous or nonaqueous medium or with LiOR ,
10 LiNH_2 , LiNR_2 , NaOR , NaNH_2 , NaNR_2 , KOR , KNH_2 or KNR_2 , wherein R is an alkyl group containing 1-4 carbon atoms, in a nonaqueous medium.

b) Salts wherein the cation is Mg^{2+} or Ca^{2+} , are prepared
15 by treating the compound of the invention with $\text{Mg}(\text{OR})_2$, $\text{Ca}(\text{OR})_2$ or CaH_2 , wherein R is an alkyl group containing 1-4 carbon atoms, in a nonaqueous solvent such as an alcohol (only for the alcoholates), e.g. ROH , or in an ether such as tetrahydrofuran.

20

Racemates obtained can be separated into the pure enantiomers. This may be done according to known methods, e.g. from racemic diastereomeric salts by means of chromatography or fractional crystallization.

25

The starting materials described in the intermediate examples may be obtained according to processes known per se.

30 For clinical use the compound of the invention is formulated into pharmaceutical formulations for oral, rectal, parenteral or other modes of administration. The pharmaceutical formulation contains the compound of the invention normally in combination with a pharmaceutically
35 acceptable carrier. The carrier may be in the form of a solid, semi-solid or liquid diluent, or a capsule. These

pharmaceutical preparations are a further object of the invention. Usually the amount of active compound is between 0.1-95% by weight of the preparation, between 0.2-20% by weight in preparations for parenteral use and
5 between 1-50% by weight in preparations for oral administration.

In the preparation of pharmaceutical formulations containing the compound of the present invention in the
10 form of dosage units for oral administration the compound selected may be mixed with a solid, powdered carrier, such as lactose, saccharose, sorbitol, mannitol, starch, amylopectin, cellulose derivatives, gelatin, or another suitable carrier, stabilizing substances such as alkaline
15 compounds e.g. carbonates, hydroxides and oxides of sodium, potassium, calcium, magnesium and the like as well as with lubricating agents such as magnesium stearate, calcium stearate, sodium stearyl fumarate and polyethylenglycol waxes. The mixture is then processed
20 into granules or pressed into tablets. Granules and tablets may be coated with an enteric coating which protects the active compound from acid catalyzed degradation as long as the dosage form remains in the stomach. The enteric coating is chosen among
25 pharmaceutically acceptable enteric-coating materials e.g. beeswax, shellac or anionic film-forming polymers such as cellulose acetate phthalate, hydroxypropyl-methylcellulose phthalate, partly methyl esterified methacrylic acid polymers and the like, if preferred in combination with a
30 suitable plasticizer. To the coating various dyes may be added in order to distinguish among tablets or granules with different active compounds or with different amounts of the active compound present.

35 Soft gelatine capsules may be prepared with capsules containing a mixture of the active compound of the

invention, vegetable oil, fat, or other suitable vehicle for soft gelatine capsules. Soft gelatine capsules may also be enteric-coated as described above. Hard gelatine capsules may contain granules or enteric-coated granules of the active compound. Hard gelatine capsules may also contain the active compound in combination with a solid powdered carrier such as lactose, saccharose, sorbitol, mannitol, potato starch, amylopectin, cellulose derivatives or gelatine. The hard gelatine capsules may be enteric-coated as described above.

Dosage units for rectal administration may be prepared in the form of suppositories which contain the active substance mixed with a neutral fat base, or they may be prepared in the form of a gelatine rectal capsule which contains the active substance in a mixture with a vegetable oil, paraffin oil or other suitable vehicle for gelatine rectal capsules, or they may be prepared in the form of a ready-made micro enema, or they may be prepared in the form of a dry micro enema formulation to be reconstituted in a suitable solvent just prior to administration.

Liquid preparation for oral administration may be prepared in the form of syrups or suspensions, e.g. solutions or suspensions containing from 0.2% to 20% by weight of the active ingredient and the remainder consisting of sugar or sugar alcohols and a mixture of ethanol, water, glycerol, propylene glycol and/or polyethylene glycol. If desired, such liquid preparations may contain colouring agents, flavouring agents, saccharine and carboxymethyl cellulose or other thickening agents. Liquid preparations for oral administration may also be prepared in the form of a dry powder to be reconstituted with a suitable solvent prior to use.

Solutions for parenteral administration may be prepared as a solution of the compound of the invention in a pharmaceutically acceptable solvent, preferably in a concentration from 0.1% to 10% by weight. These solutions
5 may also contain stabilizing agents and/or buffering agents and may be manufactured in different unit dose ampoules or vials. Solutions for parenteral administration may also be prepared as a dry preparation to be reconstituted with a suitable solvent
10 extemporaneously before use.

The typical daily dose of the active substance will depend on various factors such as for example the individual requirement of each patient, the route of
15 administration and the disease. In general, oral and parenteral dosages will be in the range of 5 to 500 mg per day of active substance.

The invention is illustrated by the following examples.
20

Example 1 Preparation of 5-fluoro-2-[[[4-cyclopropylmethoxy-2-pyridinyl)methyl]sulfinyl]-1H-benzimidazole

25 5- Fluoro-2-[[[4-cyclopropylmethoxy-2-pyridinyl)methyl]-thio]-1H-benzimidazole (1.25 g, 0.0036 mol) was dissolved in CH_2Cl_2 (40 ml). NaHCO_3 (0.6 g, 0.0072 mol) dissolved in H_2O (20 ml) was added and the mixture was cooled to +2°C. m-Chloroperbenzoic acid, 84% (0.73 g, 0.0036 mol)
30 dissolved in CH_2Cl_2 (5 ml) was added under stirring. Stirring was continued at room temperature for 15 min. The two phases were separated and NaOH (0.29 g, 0.0072 mol) dissolved in H_2O (25 ml) was added to the organic phase. The mixture was stirred, the phases were separated and the
35 H_2O phase was treated with Norite and filtered. Methylformiate (0.45 ml, 0.0073 mol) dissolved in H_2O

(5 ml) was added dropwise under stirring. After extraction with CH_2Cl_2 and drying with Na_2SO_4 the solvent was evaporated. In this way the title compound was obtained (0.93 g, 69%). NMR data for the final product is given below.

Example 2. Preparation of 5-fluoro-2-[[[4-cyclopropylmethoxy-2-pyridinyl)methyl]sulfinyl]-1H-benzimidazole, sodium salt

10 5-Fluoro-2-[[[4-cyclopropylmethoxy-2-pyridinyl)methyl]sulfinyl]-1H-benzimidazole (5 g; 14.5 mmol) dissolved in dichloromethane (100 ml) and sodium hydroxide (0.56 g; 14 mmol) dissolved in water (100 ml) were transferred to a
15 separatory funnel. The mixture was shaken to equilibrium whereupon the solvent phases were separated. The water solution was washed with dichloromethane (2 x 25 ml) and then freeze dried. The residue was recrystallized from dichloromethane/diethyl ether. Yield: 3.7 g (71 %) of the
20 title compound. NMR data is given below.

Table 1

5	Ex.	Solvent	NMR data δ ppm (500 MHz)
10	1.	CDCl ₃	0.22 (m, 2H); 0.60 (m, 2H); 1.10 (m, 1H); 3.45 (m, 1H); 3.60 (m, 1H); 4.52 (d, 1H); 4.70 (d, 1H); 6.65 (d, 1H); 6.70 (dd, 1H); 7.08 (m, 1H); 7.30-7.90 (b, 2H); 8.28 (d, 1H)
15	2.	D ₂ O δ (D ₂ O, 4.82)	0.09 (m, 2H); 0.49 (m, 2H); 0.88 (m, 1H); 2.92 (m, 1H); 3.34 (m, 1H); 4.62 (d, 1H); 4.71 (d, 1H); 6.05 (d, 1H); 6.75 (m, 1H); 7.05 (m, 1H); 7.33 (m, 1H); 7.58 (m, 1H); 8.23 (d, 1H)
20			

Preparation of synthetic intermediates

5 Example I 1. Preparation of 4-cyclopropylmethoxy-2-methylpyridine-1-oxide.

To sodium hydride (55% pure) (4.4 g, 0.1 mol) (washed with petroleum ether), cyclopropyl-methanol (50 ml) was added. Then a solution of 2-methyl-4-nitropyridine-N-oxide (6.5
10 g, 0.042 mol) in cyclopropylmethanol (30 ml) was added during about 1 h. The dark brown mixture was heated to 90°C and stirred at 90° C for about 1 h. Thereafter the cyclopropylmethanol was distilled off under reduced pressure and methylene chloride (100 ml) was added to the
15 residue. The mixture was stirred for about 30 minutes, then filtered and concentrated which gave 9.5 g of crude material.

The crude material was purified by flash chromatography on
20 silica with methylene chloride-methanol (90-10) as eluent, giving 4.0 g (53%) of pure title compound. NMR data is given below.

25 Example I 2. Preparation of 2-acetoxymethyl-4-cyclopropylmethoxypyridine.

4-cyclopropylmethoxy-2-methylpyridine-1-oxide (3.8 g 0.021 mol) was dissolved in acetic anhydride (10 ml) and was added dropwise to acetic anhydride (20 ml) (warmed to
30 90°C). After the addition the temperature was raised to 110°C and the mixture was stirred at 110°C for 1 h and then the solvent was distilled off and the crude product was used without purification. NMR data is given below.

Example I 3. Preparation of 4-cyclopropylmethoxy-2-hydroxymethylpyridine

5 To the crude 2-acetoxymethyl-4-cyclopropylmethoxy
pyridine, NaOH (100 ml 2 M) was added and the mixture was
refluxed for 2 hours. The mixture was extracted with
methylene chloride, and the phases were separated. The
organic layer was dried with Na₂SO₄, filtered and the
10 solvent was evaporated off, yielding 2.7 g of crude title
compound. NMR data is given below. The crude product was
used without any further purification.

Example I 4. Preparation of 4-cyclopropylmethoxy-2-
15 chloromethylpyridine hydrochloride

4-cyclopropylmethoxy-2-hydroxymethyl pyridine (93% pure)
(0.9 g 0.0046 mol) was dissolved in methylene chloride (10
20 ml) and cooled to 0°C. SOCl₂ (0.5 ml, 0.0069 mol) in
methylene chloride (5 ml) was added dropwise at 0°C and
the reaction mixture was stirred 15 min at room
temperature. Isopropanol (0.5 ml) was added and the
mixture was evaporated giving the desired product (0.68 g,
25 78%). NMR-data is given below.

Example I 5. Preparation of 5-fluoro-2-[[[4-
cyclopropylmethoxy-2-pyridinyl)methyl)thio]-1H-
30 benzimidazole used as starting material

To 5-fluoro-2-mercapto-1H-benzimidazole (0.88 g, 0.0051
mol) in methanol (25 ml) NaOH (0.2 g, 0.0051 mol)
35 dissolved in H₂O (1 ml) and 4-cyclopropylmethoxy-2-
chloromethyl-pyridine hydrochloride (0.91 g, 0.0046 mol)

dissolved in methanol (10 ml) were added in the given order. The mixture was heated to boiling and NaOH (0.2 g, 0.005 mol) dissolved in H₂O (1 ml) was added and the mixture was refluxed for 1 hour. After evaporation of
5 methanol, CH₂Cl₂ (75 ml) and H₂O (50 ml) were added and pH adjusted to 10. The mixture was vigorously stirred, the phases were separated, the organic phase was dried over Na₂SO₄ and evaporated giving the desired product (1.25 g, 72%). NMR data for the product is given below.

Table 2.

Ex	Solvent	NMR data δ ppm
5	I 1. CDCl ₃ (500 MHz)	0.36 (m, 2H); 0.68 (m, 2H); 1.26 (m, 1H); 2.52 (s, 3H); 3.83 (d, 2H); 6.70 (dd, 1H); 6.77 (d, 1H); 8.16 (d, 1H)
10	I 2. CDCl ₃ (500 MHz)	0.37 (m, 2H); 0.69 (m, 2H); 2.16 (s, 3H); 3.87 (d, 2H); 6.75 (dd, 1H); 6.87 (d, 1H); 8.42 (d, 1H)
15	I 3. CDCl ₃ (500 MHz)	0.36 (m, 2H); 0.67 (m, 2H); 1.27 (m, 1H); 3.86 (d, 2H); 4.69 (s, 2H); 6.72 (dd, 1H); 6.78 (d, 1H); 8.33 (d, 1H)
20	I 4. DMSO-d ₆ (300 MHz)	0.40 (m, 2H); 0.60 (m, 2H); 1.30 (m, 1H); 4.20 (d, 2H); 5.00 (s, 2H); 7.45 (dd, 1H); 7.65 (d, 1H); 8.70 (d, 1H)
25	I 5. CDCl ₃ (500 MHz)	0.36-0.39 (m, 2H); 0.67-0.71 (m, 2H); 1.27 (m, 1H); 3.89 (d, 2H); 4.29 (s, 2H); 6.81 (dd, 1H); 6.89 (d, 1H); 6.94 (m, 1H); 7.24 (dd, 1H); 7.46 (dd, 1H); 8.43 (d, 1H)

30

The best mode of carrying out the invention known at present is to use the sodium salt of the compound of the invention, thus the compound described in Example 2.

Pharmaceutical preparations containing the compound of the invention as active ingredient are illustrated in the following formulations.

5 Syrup

A syrup containing 1% (weight per volume) of active substance was prepared from the following ingredients:

	Compound according to Example 1	1.0 g
10	Sugar, powder	30.0 g
	Saccharine	0.6 g
	Glycerol	5.0 g
	Flavouring agent	0.05 g
	Ethanol 96%	5.0 g
15	Distilled water q.s. to a final volume of	100 ml

Sugar and saccharine were dissolved in 60 g of warm water. After cooling the active compound was added to the sugar solution and glycerol and a solution of flavouring agents
20 dissolved in ethanol were added. The mixture was diluted with water to a final volume of 100 ml.

Enteric-coated tablets

An enteric coated tablet containing 50 mg of active
25 compound was prepared from the following ingredients:

	I Compound according to Example 1	500 g
	as Mg salt	
30	Lactose	700 g
	Methyl cellulose	6 g
	Polyvinylpyrrolidone cross-linked	50 g
	Magnesium stearate	15 g
	Sodium carbonate	6 g
35	Distilled water	q.s.

II	Cellulose acetate phthalate	200 g
	Cetyl alcohol	15 g
	Isopropanol	2000 g
5	Methylene chloride	2000 g

I Compound according to example 1, powder, was mixed with lactose and granulated with a water solution of methyl cellulose and sodium carbonate. The wet mass was forced through a sieve and the granulate dried in an oven. After drying the granulate was mixed with polyvinylpyrrolidone and magnesium stearate. The dry mixture was pressed into tablet cores (10 000 tablets), each tablet containing 50 mg of active substance, in a tabletting machine using 7 mm diameter punches.

II A solution of cellulose acetate phthalate and cetyl alcohol in isopropanol/methylene chloride was sprayed onto the tablets I in an Accela Cota[®], Manesty coating equipment. A final tablet weight of 110 mg was obtained.

Solution for intravenous administration

A parenteral formulation for intravenous use, containing 4 mg of active compound per ml, was prepared from the following ingredients:

Compound according to Example 2

30	Sterile water to a final volume of	4 g 1000 ml
----	------------------------------------	----------------

The active compound was dissolved in water to a final volume of 1000 ml. The solution was filtered through a 0.22 μ m filter and immediately dispensed into 10 ml sterile ampoules. The ampoules were sealed.

Capsules

Capsules containing 30 mg of active compound were prepared from the following ingredients:

5

Compound according to Example 1	300 g
Lactose	700 g
Microcrystalline cellulose	40 g
Hydroxypropyl cellulose low-substituted	62 g
10 Disodium hydrogen phosphate	2 g
Purified water	q.s.

The active compound was mixed with the dry ingredients and granulated with a solution of disodium hydrogen
15 phosphate. The wet mass was forced through an extruder and spheronized and dried in a fluidized bed dryer.

500 g of the pellets above were first coated with a solution of hydroxypropyl methylcellulose, 30 g, in water,
20 750 g, using a fluidized bed coater. After drying, the pellets were coated with a second coating as given below:

Coating solution:

25 Hydroxypropyl methylcellulose phthalate	70 g
Cetyl alcohol	4 g
Acetone	200 g
Ethanol	600 g

30 The final coated pellets were filled into capsules.

Suppositories

Suppositories were prepared from the following ingredients using a welding procedure. Each suppository contained 40 mg of active compound.

Compound according to Example 1	4 g
Witepsol H-15	180 g

The active compound was homogenously mixed with Witepsol H-15 at a temperature of 41°C. The molten mass was volume filled into pre-fabricated suppository packages to a net weight of 1.84 g. After cooling the packages were heat sealed. Each suppository contained 40 mg of active compound.

Biological Effects

Bioavailability

Choice of Species for Testing.

The results from tests on two different animal species, rat and dog, vary in regard to measured level of bioavailability for the same compound. We believe that the rat is the more relevant species for bioavailability testing. This is based on our belief that liver metabolism has the most predominant impact upon bioavailability, and that the liver metabolic pattern in man for this type of compounds is quite similar to that of

the male rat (more so than of the female rat and the dog). Moreover, test results of bioavailability in the male rat will tend to give a broader "spread" compared with the test results in the dog, and thus the male rat model will give more clear differences in bioavailability between different compounds. Stated in another way, the bioavailability as tested in the male rat can be expected to give a better estimate of the relative differences in man between different test compounds compared with the test results obtained when using the same compounds in the dog.

Assessment of Bioavailability.

Bioavailability is assessed by calculating the quotient between the area under plasma concentration (AUC) curve following intraduodenal (id) administration and intravenous (iv) administration from the rat or the dog. Low, therapeutically relevant doses, were used. This method is scientifically recognized as valid for assessing bioavailability (see for instance: M. Rowland and T.N. Tozer, Clinical Pharmacokinetics, 2nd ed., Lea & Febiger, London 1989, p 42). The data from both the rat and the dog are provided in Table 3.

Rough Screening Model.

Since the bioavailability model described above is time and labour intensive, and requires a large number of plasma analyses, also a rough screening model, based on relative potencies to inhibit acid secretion, has been used (see for instance: A. Goth, Medical Pharmacology, 7th ed., C.V. Mosby Company, Saint Louis 1974, p 19). Thus, the ratio (called "Bioavailability" in Table 3) between the ED_{50} at intravenous administration and the ED_{50} at

intraduodenal administration was calculated. Also these data are provided in Table 3.

Potency

5 The potency for inhibition of acid secretion has been measured in the male rat and the dog, both intravenously and intraduodenally. When it comes to relevance of the animal test data for potency of a given compound in man
10 for the present type of compounds, it is believed that potency in man will correspond to a level somewhere between what is measured in the male rat and what is measured in the dog. Potency data from the two animal species are given in Table 3.

15

Biological Tests

Inhibition of Gastric Acid Secretion in the Conscious Male Rat.

20

Male rats of the Sprague-Dawley strain were used. They were equipped with cannulated fistulae in the stomach (lumen) and the upper part of the duodenum, for collection of gastric secretions and administration of test
25 substances, respectively. A fourteen days recovery period after surgery was allowed before testing commenced.

Before secretory tests, the animals were deprived of food but not water for 20 h. The stomach was repeatedly washed
30 through the gastric cannula, and 6 ml of Ringer-Glucose given s.c. Acid secretion was stimulated with a infusion during 3.5 h (1.2 ml/h, s.c.) of pentagastrin and carbachol (20 and 110 nmol/kg h, respectively), during which time gastric secretions were collected in 30-min
35 fractions. Test substances or vehicle were given iv or id at 90 min after starting the stimulation, in a volume of 1

ml/kg. Gastric juice samples were titrated to pH 7.0 with NaOH, 0.1 mol/L, and acid output calculated as the product of titrant volume and concentration. Further calculations were based on group mean responses from 4-5 rats. The acid output during the periods after administration of test substances or vehicle were expressed as fractional responses, setting the acid output in the 30-min period preceding administration to 1.0. Percentage inhibition was calculated from the fractional responses elicited by test compound and vehicle. ED₅₀ values were obtained from graphical interpolation on log dose-response curves, or estimated from single-dose experiments assuming a similar slope for all dose-response curves. An estimation of the bioavailability was obtained by calculating the ratio ED_{50iv}/ED_{50id}. The results reported are based on gastric acid secretion during the second hour after drug/vehicle administration.

Bioavailability in the Male Rat.

Male adult rats of the Sprague-Dawley strain were used. One day, prior to the experiments, all rats were prepared by cannulation of the left carotid artery under anaesthesia. The rats used for the intravenous experiments, were also cannulated in the jugular vein. (Ref. V Popovic and P Popovic, J Appl Physiol 1960;15,727-728). The rats used for the intraduodenal experiments, were also cannulated in the upper part of the duodenum. The cannulas were exteriorized at the nape of the neck. The rats were housed individually after surgery and were deprived of food, but not water, before administration of the test substances. The same dose (4 µmol/kg) were given iv and id as a bolus for about one minute (2 ml/kg). Blood samples (0.1-0.4 g) were drawn repeatedly from the carotid artery at intervals up to 4 hours after given

dose. The samples were frozen as soon as possible until analysis of the test compound.

The area under the blood concentration vs time curve, AUC, was determined by the linear trapezoidal rule and extrapolated to infinity by dividing the last determined blood concentration by the elimination rate constant in the terminal phase. The systemic bioavailability (F%) following intraduodenal administration was calculated as

10

$$F(\%) = \frac{AUC_{id}}{AUC_{iv}} \times 100$$

15 Inhibition of Gastric Acid Secretion and Bioavailability in the Conscious Dog.

Harrier dogs of either sex were used. They were equipped with a duodenal fistula for the administration of test compounds or vehicle and a cannulated ventricular fistula for the collection of gastric secretions.

Before secretory tests the animals were fasted for about 18 h but water was freely allowed. Gastric acid secretion was stimulated by a 4 h infusion of histamine dihydrochloride (12 ml/h) at a dose producing about 80% of the individual maximal secretory response, and gastric juice collected in consecutive 30-min fractions. Test substance or vehicle was given id or iv 1 h after starting the histamine infusion, in a volume of 0.5 ml/kg body weight. The acidity of the gastric juice samples were determined by titration to pH 7.0, and the acid output calculated. The acid output in the collection periods after administration of test substance or vehicle were expressed as fractional responses, setting the acid output in the fraction preceding administration to 1.0.

Percentage inhibition was calculated from fractional responses elicited by test compound and vehicle. ED_{50} -values were obtained by graphical interpolation on log dose - response curves, or estimated from single-dose experiments under the assumption of the same slope of the dose-response curve for all test compounds. All results reported are based on acid output 2 h after dosing.

Blood samples for the analysis of test compound concentration in plasma were taken at intervals up to 3 h after dosing. Plasma was separated and frozen within 30 min after collection. AUC (area under the plasma concentration - time curve), extrapolated to infinite time, was calculated by the linear trapezoidal rule. The systemic bioavailability (F%) after id administration was calculated as $100 \times (AUC_{id}/AUC_{iv})$.

Chemical Stability

The chemical stability of various compounds of the invention has been followed kinetically at low concentration at 37°C in aqueous buffer solution at different pH values. The results in Table 3 show the half life ($t_{1/2}$) at pH 7, that is the time period after which half the amount of the original compound remains unchanged.

Results of biological and stability tests

Table 3 gives a summary of the test data available for the compound of the invention and a structurally closely related compound in the prior art, called Ref. in Table 3, namely 5-fluoro-2-[[[(4-isopropoxy-2-pyridinyl)-methyl]sulfinyl]-1H-benzimidazole described in US 4 727 150. As can be seen from Table 3 the compound according to the invention has a high bioavailability (F =

82% in the rat), high potency ($ED_{50iv} = 1.2 \mu\text{mol/kg}$, $ED_{50id} = 2.2 \mu\text{mol/kg}$ in the rat) and a high chemical stability ($t_{1/2} = 23 \text{ h}$). Moreover, considering the most distinguishing property for the compound of the invention, 5 the bioavailability, the compound of the invention has a much higher value (82% vs 31%) compared to that of the Ref. compound, and is better in the other properties as well ($ED_{50iv} = 1.8 \mu\text{mol/kg}$, $ED_{50id} = 4.0 \mu\text{mol/kg}$ and $t_{1/2} = 14 \text{ h}$ for the Ref compound).

Table 3, Biological Test Data and Stability Data

Test compound Example no.	Inhibition of acid secretion		"Bioavailability" measured by the Rough Screening Model		Bioavailability measured by the AUC-method F%		Chemical stability at pH 7
	Dog ED ₅₀ (μmol/kg)		Rat ED ₅₀ iv/ ED ₅₀ id (%)				
	Route of adm. iv	adm. id	Route of adm. iv	adm. id	Dog	Rat	
1	1)	1.0	1.2	2.2	80	82	23
Ref.	n.t.	2)	1.8	4.0	n.t.	31	14

n.t. = not tested

1) Dog 1 1 $\mu\text{mol/kg}$ gave 35% inhibition
 Dog 2 1 " gave no effect
 Dog 3 2 " gave 98% inhibition.

2) Dog 4 3 $\mu\text{mol/kg}$ gave 95% inhibition
 Dog 5 3 " gave 98% inhibition

Thus no ED₅₀ value could be estimated.Thus no ED₅₀ value could be estimated.

CLAIMS

1. 5-Fluoro-2-[[[4-cyclopropylmethoxy-2-pyridinyl)methyl]-
5 sulfinyl]-1H-benzimidazole (compound I) and physiologically
acceptable salts thereof, as well as its optical enantiomers.
2. The sodium salt of the compound according to claim 1.
- 10 3. The magnesium salt of the compound according to claim 1.
4. A pharmaceutical composition containing as active
ingredient the compound according to claim 1.
- 15 5. A compound as defined in claim 1 for use in therapy.
6. A compound as defined in claim 1 for use in inhibiting
gastric acid secretion in mammals including man.
- 20 7. A compound as defined in claim 1 for use in the
treatment of gastrointestinal inflammatory diseases in
mammals including man.
8. Use of a compound according to claim 1 for the
25 manufacture of a medicament for inhibiting gastric acid
secretion in mammals including man.
9. Use of a compound according to claim 1 for the
manufacture of a medicament for the treatment of
30 gastrointestinal inflammatory diseases in mammals including
man.
10. A process for the preparation of a compound according to
claim 1, by oxidizing 5-fluoro-2-[[[4-cyclopropylmethoxy-2-
35 pyridinyl)methyl]thio]-1H-benzimidazole whereupon, compound

I thus obtained if desired, is converted to a salt or into a pure optical isomer.

11. 4-cyclopropylmethoxy-2-methylpyridine-1-oxide.
- 5 12. 2-acetoxymethyl-4-cyclopropylmethoxypyridine.
13. 4-cyclopropylmethoxy-2-hydroxymethylpyridine.
- 10 14. 4-cyclopropylmethoxy-2-chloromethylpyridine
hydrochloride.
- 15 15. 5-fluoro-2-[[(4-cyclopropylmethoxy-2-pyridinyl)methyl]-
thio]-1H-benzimidazole.

INTERNATIONAL SEARCH REPORT

International Application No **PCT/SE 89/00740**

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) *		
According to International Patent Classification (IPC) or to both National Classification and IPC		
IPC5: C 07 D 401/12, 213/68, 213/89, A 61 K 31/44		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
IPC5	C 07 D	
Documentation Searched other than Minimum Documentation = the extent that such Documents are included in the Fields Searched *		
SE,DK,FI,NO classes as above		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category ⁸	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
A	EP, A1, 0175464 (TAKEDA CHEMICAL INDUSTRIES LTD) 26 March 1986, see the whole document ----- -----	1-15
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"Z" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
2nd March 1990		1990 -03- 14
International Searching Authority		Signature of Authorized Officer
SWEDISH PATENT OFFICE		Göran Karlsson

PCT/SE 89/00740

五、

一、

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER: _____**

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem-Mailbox.